


# EP 200ng

## DATASHEET

Version: 01-03-2020

EP 200ng is a complementary feed produced from Non-GMO soybean meal fermented with lactic acid bacteria. EP200ng contains fermentation metabolites, including lactic acid, and gut flora stabilizing lactic acid bacteria. The fermentation process makes the product highly digestible. EP200ng is suitable for young and vulnerable animals and helps maintaining a healthy and functional gut.



Better gut  
health for  
your animals

## NUTRIENTS

DM	895 g/kg
Crude Protein	505 g/kg
Crude Fat	22 g/kg
Crude Fibre	34 g/kg
Crude Ash	64 g/kg
Starch	41 g/kg
Sugars	14 g/kg

## ENERGY

ME Pigs	14.7 MJ/kg
ME Broilers	10.0 MJ/kg
ME Layers	10.4 MJ/kg

## MINERALS

Calcium	2.6 g/kg
Phosphorus	7.3 g/kg
Dig. Phosphorus	4.7 g/kg
Sodium	0.2 g/kg
Potassium	21.7 g/kg
Chloride	0.3 g/kg

## LACTIC ACID AND BACTERIA

Lactic acid	73 g/kg
Lactic acid bacteria*	> 10 <sup>6</sup> cfu/g

\* Enterococcus faecium (NCIMB 10415)

AMINO ACIDS	Gross	SID Pigs	SID Poultry	
Lys	27.9	24.0	24.3	g/kg
Meth	7.0	6.5	6.6	g/kg
M+C	14.6	12.8	13.6	g/kg
Thre	19.8	17.6	18.6	g/kg
Tryp	6.5	5.9	6.2	g/kg
Ile	23.1	21.0	21.7	g/kg
Arg	35.4	33.6	34.3	g/kg
His	13.6	12.2	12.9	g/kg
Leu	38.9	35.4	37.0	g/kg
Val	24.2	21.8	22.3	g/kg
Phen	26.5	24.4	25.2	g/kg
Tyr	18.7	17.0	17.6	g/kg

## ANTINUTRITIONAL SUBSTANCES

TIA	< 0.5	mg/g
$\beta$ -conglycinine	< 0.5	mg/g
Lectins	< 0.05	mg/g
Raffinose and Stachyose	< 0.5	%

EP200 is a fermented complementary feed composed out of natural Non-GMO raw materials. The figures are based on analyses representing an average. Some variance in chemical composition may occur.

**Physical characteristic appearance:** Yellowish light brown meal product with a pleasant flavor of fermentation.

**Package size:** 1.000 kg in big bags

**Storage & Shelf life:** 12 months from production date when stored at 20°C or below. Shelf life will be shorter if stored at a higher temperature.

### Instructions for use\*:

	Period	EP200
<b>PIGS</b>		
<b>Piglets</b>	Before and after weaning	8-15%
	12-20 kg	1-8%
<b>POULTRY</b>		
<b>Broilers</b>	Until day 22	5-10%
	After day 22	2-5%
<b>AQUA</b>		
<b>Salmons</b>		15-25%

\*Percent of dry feed.

The product contains 7% lactic acid. Remember to optimize and reduce added acidifiers.