


# EP 200

## DATASHEET

Version: 01-03-2020

EP 200 is a complementary feed produced from soybean meal fermented with lactic acid bacteria. EP200 contains fermentation metabolites, including lactic acid, and gut flora stabilizing lactic acid bacteria. The fermentation process makes the product highly digestible. EP200 is suitable for young and vulnerable animals and helps maintaining a healthy and functional gut.



Better gut  
health for  
your animals

## NUTRIENTS

DM	890 g/kg
Crude Protein	495 g/kg
Crude Fat	23 g/kg
Crude Fibre	34 g/kg
Crude Ash	65 g/kg
Starch	38 g/kg
Sugars	26 g/kg

## ENERGY

ME Pigs	14.8 MJ/kg
ME Broilers	10.0 MJ/kg
ME Layers	10.4 MJ/kg

## MINERALS

Calcium	3.8 g/kg
Phosphorus	6.7 g/kg
Dig. Phosphorus	4.3 g/kg
Sodium	0.2 g/kg
Potassium	21.6 g/kg
Chloride	0.4 g/kg

## LACTIC ACID AND BACTERIA

Lactic acid	74 g/kg
Lactic acid bacteria*	> 10 <sup>6</sup> cfu/g

\* Enterococcus faecium (NCIMB 10415)

AMINO ACIDS	Gross	SID Pigs	SID Poultry	
Lys	29.0	24.9	25.2	g/kg
Meth	6.9	6.4	6.5	g/kg
M+C	13.8	12.1	12.8	g/kg
Thre	20.7	18.4	19.4	g/kg
Tryp	6.7	6.1	6.5	g/kg
Ile	21.9	19.9	20.6	g/kg
Arg	34.0	32.3	33.0	g/kg
His	12.8	11.5	12.2	g/kg
Leu	37.9	34.5	36.0	g/kg
Val	23.0	20.7	21.1	g/kg
Phen	25.3	23.2	24.0	g/kg
Tyr	18.1	16.5	17.0	g/kg

## ANTINUTRITIONAL SUBSTANCES

TIA	< 0.5	mg/g
$\beta$ -conglycinine	< 0.5	mg/g
Lectins	< 0.05	mg/g
Raffinose and Stachyose	<0.5	%

EP200 is a fermented complementary feed composed out of natural raw materials. The figures are based on analyses representing an average. Some variance in chemical composition may occur. This product contains genetically modified soybean meal.

**Physical characteristic appearance:** Yellowish light brown meal product with a pleasant flavor of fermentation.

**Package size:** Bulk or 1.000 kg in big bags.

**Storage & Shelf life of nutrients:** 12 months from production date when stored at 20°C or below. Shelf life will be shorter if stored at a higher temperature.

### Instructions for use\*:

	Period	EP200
<b>PIGS</b>		
<b>Piglets</b>	Before and after weaning	8-15%
	12-20 kg	1-8%
<b>POULTRY</b>		
<b>Broilers</b>	Until day 22	5-10%
	After day 22	2-5%
<b>AQUA</b>		
<b>Salmons</b>		15-25%

\*Percent of dry feed.

The product contains 7% lactic acid. Remember to optimize and reduce added acidifiers.